Opportunities for Merging Landsat and Sentinel-2 Data Streams

Jeffrey Masek, NASA GSFC John Dwyer, USGS EROS Garik Gutman, NASA HQ


Dec 12, 2012

Moderate-Resolution Time Series


Since the opening of the USGS Landsat archive, there has been increased interest in *intra-annual* time series applications at 30m resolution

- Agricultural monitoring
- Vegetation biophysics (LAI, fPAR, productivity)
- Phenology and climate linkages
- WELD data products (SDSU/Roy)

Example: New England forest phenology from multi-annual Landsat observations, courtesy Mark Friedl (BU)



Proposed Sentinel-2 / Landsat Architecture




Activities

- 1. Develop algorithms to make Sentinel-2 and Landsat/LDCM data compatible
- Support USGS in securing access to Sentinel data for US investigators
- 3. Pursue prototyping GEO-GLAM agricultural monitoring (crop area, type, condition)
 - Collaborate with USDA, USGS, JRC MARS
 - Phased approach: test sites -> large-area

1. Merged Radiometric Record

- Collaboration with UMD, CNES/CESBIO, GSFC, ARC, USGS
 - First meeting held Oct 25-26 2012 @ GSFC
- Goal: "seamless" surface reflectance product with <5 day repeat to support GEO-GLAM
 - Three products identified: Landsat SR (LSR), Sentinel SR (SSR), and Merged SR (MSR)
- Radiometric Adjustments
 - Cross-calibration
 - Atmospheric correction (6S, MACC)
 - BRDF correction to nadir & constant SZA
 - Bandpass correction
 - Regridding and compositing (TBD)
- Initial prototyping using SPOT-4 and L7 data (Spring 2013)

Merged Radiometric Record



2. Sentinel-2 Data Access: Background

- Sentinel-2 data will be transmitted via X-band to four ground stations
 - Alaska, Svalbard, Matera, and Maspalomas
 - No processing capabilities will be hosted at these ground stations
- Mission data will be sent to Processing and Archive Centers (PACs)
 - United Kingdom
 - Spain
- PACs will process and distribute Level-1b (radiometrically corrected) and Level-1c (precision/terrain corrected) products

ESA Collaboration with NASA and USGS

- NASA will be the principal U.S. agency to work with ESA on Sentinel-1 and -3 collaboration
- USGS will be the principal agency to work with ESA on Sentinel-2 collaboration
- NASA and the USGS will work seamlessly across agencies to address topics of mutual interest
 - Establish mirror archives
 - Science and applications support (e.g. agriculture)

Sentinel-2 Mirror Archives

- The USGS is interested in hosting mirror archives of Sentinel-2 data
 - Initially will focus on Level-1c products (100 Km by 100 Km granules) as soon as they are made available from the PACs
 - Ensure synchronization of U.S. and ESA inventories
 - US would serve as a "hot back-up"
 - May investigate Level-1b products
 - To enable generation of larger Level-1c granules conforming to Landsat WRS-2 scenes or WELD tiles
 - To enable the generation of high-level geophysical and biophysical parameters
- USGS also investigating direct downlink options for S2

Data Access Challenges

- Sentinel-2 Data Policy and Operations Budget still unclear
 - Announcement at EU US Space Dialogue that all of the Sentinel data will be free and open with no copyright issues.
- Access to science processing software from ESA or Vendors is to be resolved
 - May be limited to binary executables rather than source code
 - Cost and terms of software licenses unknown
- Need to establish network capability between Europe and U.S.
 - Martha Maiden has requested the ESDIS project to investigate potential solutions
- Support of near real-time applications in U.S. or North America may not be feasible

Conclusions

- Combining Sentinel-2 and Landsat observations can provide global, <5 day coverage
- USGS and NASA are engaged in establishing a framework for...
 - US access to Sentinel-2 data
 - Algorithm and processing approaches for creating a merged data stream
 - Improved LCLUC and agricultural monitoring applications via merged data stream
- LST engagement highly desired

TABLE 1: TASKS PLAN

10 CESBIO meeting (see section 7) Meeting 07/2013 CESBIO					
1 Take-five and insure the Landsat-7 and LDCM acquisition Make a choice on the spatial resolution of the MSR product by (ii) analyzing field size distribution based on geowiki data (see D2_P2 slide 46) (i) Analyzing spatial consistency of Landsat-7 and LDCM based on very high spatial resolution data (<5m). Make a choice on the temporal and spectral characteristics of the MSR product Amake a choice on the temporal and spectral characteristics of the MSR product Make a choice on the projection systems, the tiling system of MSR Design a method to retrieve high spatial resolution BRDF coefficient based on coarse spatial resolution BRDF coefficient (MODIS) desegregation. Product definition O3/2013 MAKE D O3/2013 O3/2013 O3/2013 O3/2013 O3/2013 UMD and/or NASA-GSFC (Claverie / Vermote / Masek) UMD and/or NASA-GSFC (Claverie / Vermote / Masek) Product definition O3/2013 O3/2013	#	Tasks	category	schedule	responsibility
product by (ii) analyzing field size distribution based on geowiki data (see DZ_P2 slide 46) (i) Analyzing spatial consistency of Landsat-7 and LDCM based on very high spatial resolution data (<5m). Make a choice on the temporal and spectral characteristics of the MSR product Make a choice on the projection systems, the tiling system of MSR Design a method to retrieve high spatial resolution BRDF coefficient (MODIS) desegregation. Design a Band-pass adjustment method Design a temporal fusion method (WELD-like) Design the algorithm chain by concatenating atmospheric correction, BRDF correc	1	Take-five and insure the Landsat-7 and LDCM	SPOT-4 T5	11/2012	l '
Make a choice on the temporal and spectral characteristics of the MSR product Make a choice on the projection systems, the tiling system of MSR Make a choice on the projection systems, the tiling system of MSR Design a method to retrieve high spatial resolution BRDF coefficient based on coarse spatial resolution BRDF coefficient (MODIS) desegregation. Make a choice on the projection systems, the tiling system of MSR Product definition O3/2013 GSFC (Claverie / Vermote / Masek) UMD and/or NASA-GSFC (Claverie / Vermote) Wermote) UMD and/or NASA-GSFC (Claverie / Vermote) UMD and/or NASA-GSFC (Claverie / Vermote) Wermote) UMD and/or NASA-GSFC (Claverie / Vermote) UMD and/or NASA-GSFC (Claverie / Vermote) UMD and/or NASA-GSFC (Claverie / Vermote / Masek) Design a temporal fusion method (WELD-like) Product definition O3/2013 GSFC (Claverie / Vermote) UMD and/or NASA-GSFC (Claverie / Vermote / Masek) UMD and/or NASA-GSFC (Claverie / Vermote / Masek) Design the algorithm chain by concatenating atmospheric correction, BRDF correction, Band-pass adjustment and temporal fusion methods. Create a common dataset including data from Landsat-7 ETM+, LDCM OLI and SPOT-4 Take-five HRVIR. Design method (SEBIO meeting (see section 7) Meeting O7/2013 UMD, NASA-GSFC/ARGESBIO	2	product by (ii) analyzing field size distribution based on geowiki data (see D2_P2 slide 46) (i) Analyzing spatial consistency of Landsat-7 and LDCM based on very high spatial resolution data		03/2013	GSFC (Whitcraft /
Make a choice on the projection systems, the tiling system of MSR Design a method to retrieve high spatial resolution BRDF coefficient based on coarse spatial resolution BRDF coefficient (MODIS) desegregation. R & D O3/2013 GSFC (Claverie / Vermote / Masek) UMD and/or NASA-GSFC (Claverie / Vermote) Design a Band-pass adjustment method R & D O4/2013 O5/2013 UMD and/or NASA-GSFC (Claverie / Vermote) Design a temporal fusion method (WELD-like) Design the algorithm chain by concatenating atmospheric correction, BRDF correction, Band-pass adjustment and temporal fusion methods. Create a common dataset including data from Landsat-7 ETM+, LDCM OLI and SPOT-4 Take-five HRVIR. Design meeting (see section 7) Meeting O7/2013 UMD, NASA-GSFC (CESBIO)	3	• • •		03/2013	GSFC (Claverie /
BRDF coefficient based on coarse spatial resolution BRDF coefficient (MODIS) desegregation. R & D 03/2013 GSFC (Claverie / Vermote) UMD and/or NASA-GSFC (Claverie / Vermote) UMD and/or NASA-GSFC (Claverie / Vermote) R & D 05/2013 UMD and/or NASA-GSFC (Claverie / Vermote) UMD and/or NASA-GSFC (Claverie / Vermote) Besign a temporal fusion method (WELD-like) Design the algorithm chain by concatenating atmospheric correction, BRDF correction, Bandpass adjustment and temporal fusion methods. Create a common dataset including data from Landsat-7 ETM+, LDCM OLI and SPOT-4 Take-five HRVIR. Meeting 07/2013 UMD, NASA-GSFC/ARGESBIO	4	, , , , ,		03/2013	GSFC (Claverie /
Design a Band-pass adjustment method R & D 04/2013 GSFC (Claverie / Vermote) UMD and/or NASA-GSFC (Claverie / Vermote) R & D 05/2013 GSFC (Claverie / Vermote) R & D 05/2013 O5/2013 GSFC (Claverie / Vermote / Masek) R & D 06/2013 O6/2013 O6/2013 O6/2013 O6/2013 O7/2013	5	BRDF coefficient based on coarse spatial resolution	R & D	03/2013	GSFC (Claverie /
Design a temporal fusion method (WELD-like) R & D 05/2013 GSFC (Claverie / Vermote / Masek) Design the algorithm chain by concatenating atmospheric correction, BRDF correction, Bandpass adjustment and temporal fusion methods. Create a common dataset including data from Landsat-7 ETM+, LDCM OLI and SPOT-4 Take-five HRVIR. Design the elgorithm chain by concatenating R & D 06/2013 GSFC (Claverie / Vermote / Masek) Vermote / Masek) Adata handling 07/2013 NASA-GSFC UMD, NASA-GSFC/ARG CESBIO meeting (see section 7) Meeting 07/2013	6	Design a Band-pass adjustment method	R & D	04/2013	GSFC (Claverie /
8 atmospheric correction, BRDF correction, Bandpass adjustment and temporal fusion methods. Create a common dataset including data from Landsat-7 ETM+, LDCM OLI and SPOT-4 Take-five HRVIR. O6/2013 GSFC (Claverie / Vermote / Masek) O7/2013 NASA-GSFC Wheeting O7/2013 UMD, NASA-GSFC/ARG CESBIO	7	Design a temporal fusion method (WELD-like)	R & D	05/2013	GSFC (Claverie /
9 Landsat-7 ETM+, LDCM OLI and SPOT-4 Take-five HRVIR. 07/2013 NASA-GSFC 10 CESBIO meeting (see section 7) Meeting 07/2013 UMD, NASA-GSFC/ARGCESBIO	8	atmospheric correction, BRDF correction, Band- pass adjustment and temporal fusion methods.	R & D	06/2013	GSFC (Claverie /
10 CESBIO meeting (see section 7) Meeting 07/2013 CESBIO	9	Landsat-7 ETM+, LDCM OLI and SPOT-4 Take-five	data handling	07/2013	NASA-GSFC
11 Apply MACC atmospheric correction and cloud and Prototyping 08/2013 CESBIO (Hagolle)	10		Meeting	07/2013	UMD, NASA-GSFC/ARC, CESBIO
cloud shadow masking methods on the previously	11		Prototyping	08/2013	CESBIO (Hagolle)